

DOCUMENT RESUME

ED 160 473

SE 025 174

AUTHOR Wamsley, Mary Ann, Ed.; Vermeire, Donna M., Ed.
 TITLE Ornamental and Turfgrass Pest Control. Sale
 Publication 4074.
 INSTITUTION California Univ., Berkeley. Extension Div.; North
 Carolina State Univ., Raleigh.
 SPONS AGENCY Environmental Protection Agency, Washington, D.C.
 Office of Pesticide Programs.
 PUB DATE Jan 77
 CONTRACT EPA-68-01-2903
 NOTE 15p.; For related documents, see SE 025 173 and SE
 024 886-891; Contains occasional light and broken
 type

EDRS PRICE MF-\$0.83 HC-\$1.67 Plus Postage.
 DESCRIPTORS Agricultural Chemical Occupations; *Agricultural
 Education; *Disease Control; Environment; *Ornamental
 Horticulture; *Pesticides; Plant Growth; Plant
 Pathology; Science Education; *Turf Management

ABSTRACT

This guide gives information for recognition and control of ornamental and turf pests. Included are disease agents, insects and mites, weeds, and vertebrates. Symptoms and causes of phytotoxicity are given, and a discussion is presented of environmental concerns. Application methods and area measurement are also discussed. (BB)

 * Reproductions supplied by EDRS are the best that can be made *
 * from the original document. *

ED 160473

U.S. DEPARTMENT OF HEALTH,
EDUCATION & WELFARE
NATIONAL INSTITUTE OF
EDUCATION

THIS DOCUMENT HAS BEEN REPRODUCED EXACTLY AS RECEIVED FROM THE PERSON OR ORGANIZATION ORIGINATING IT. POINTS OF VIEW OR OPINIONS STATED DO NOT NECESSARILY REPRESENT OFFICIAL NATIONAL INSTITUTE OF EDUCATION POSITION OR POLICY.

ORNAMENTAL AND TURFGRASS PEST CONTROL

Division of Agricultural Sciences
UNIVERSITY OF CALIFORNIA

SALE PUBLICATION
4074

PRINTED JANUARY 1977

SE 025 174

The University of California's Cooperative Extension programs are available to all, without regard to race, color, or national origin. Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. James B. Kendrick, Jr., Director, Cooperative Extension, University of California.

10m-1/77-VL/ME

TABLE OF CONTENTS

	Page
Acknowledgments	3
Preface	3
Ornamentals	4
Introduction	4
Disease Agents	4
Weeds	5
Insects and Mites	6
Vertebrate Pests	7
Turfgrass	7
Introduction	7
Disease Agents	8
Weeds	9
Insects	10
Vertebrate Pests	11
Phytotoxicity	11
Environmental Concerns	11
Protecting Animals and People	12
Application	12
Area Measurements	12
Weights and Measures	13

ACKNOWLEDGMENTS

This guide has been developed by North Carolina State University under U.S. Environmental Protection Agency (EPA) contract number 68-01-2903. This contract was issued by the Training Branch, Operations Division, Office of Pesticide Programs, EPA. The leader of this group effort was Robert L. Robertson, North Carolina State University. Editors were Mary Ann Wamsley, EPA, and Donna M. Vermeire, North Carolina State University.

Contributors were:

John F. Ahrens, Connecticut Agricultural Experiment Station

Jack D. Butler, Colorado State University

Houston B. Couch, Virginia Polytechnic Institute and State University

Douglas Gaydon, Environmental Protection Agency, Region IV, Atlanta, Georgia

William M. Hoffman, Environmental Protection Agency, Washington, D.C.

Palmer Maples, Jr., Golf Course Superintendents Association of America, Atlanta, Georgia

Richard L. Miller, The Ohio State University

PREFACE

Federal regulations establish general and specific standards that you must meet before you can use certain pesticides. University of California Cooperative Extension offices can provide material which you may study to help you meet the *general* standards. This guide contains basic information to help you meet the *specific* standards for applicators who are engaged in ornamental and turfgrass pest control.

This guide will give you information about:

- recognition and control of ornamental pests,
- recognition and control of turfgrass pests, and
- environmental concerns for ornamental and turfgrass pest control.

This version of the ornamental and turfgrass pest control guide, designed to assist pesticide applicators in California, was edited by M. W. Schimann, Pesticide Applicator Training Coordinator, University of California Cooperative Extension. Contributors to the California version of the guide were: Clyde Elmore, University of California, Davis; Carl Koehler, University of California, Berkeley; and Arthur McCain, University of California, Berkeley.

This publication provides essential information needed to meet the Environmental Protection Agency standards on Ornamental and Turfgrass Pest Control, and to help prepare you to take the written examination for certification. It is not intended to provide you with all the knowledge you need for effective ornamental and turfgrass pest control. Additional information can be obtained from Cooperative Extension, University of California, and from your local county agricultural commissioner.

ORNAMENTALS

INTRODUCTION

Some plant damage is caused by living pests, including:

- disease agents,
- weeds,
- insects and mites, and
- vertebrate animals.

Other causes of plant problems are:

- too little, too much, or imbalanced fertilizer,
- pesticide injury,
- improper planting and pruning,
- root girdling,
- soil conditions (such as improper drainage, compaction),
- mechanical damage (by such things as earth-moving equipment, mowers, and hand tools),
- pollution damage, and
- natural aging of plants (often mistaken for damage caused by insects and diseases).

You must diagnose the problem before using control methods. Can you find an insect or recognize the symptoms of a disease? Pinpointing causes of plant damage usually requires close observation.

DISEASE AGENTS

Fungi, bacteria, viruses, nematodes, mycoplasmas, and parasitic plants cause diseases of landscape plants. Most common diseases are caused by fungi. The environment is of major importance to the development of disease in woody plants. For example:

- A sudden drop in temperature in the fall or early winter increases the susceptibility of plants to cankers caused by fungi.
- Waterlogging of the soil contributes to the development of certain root rots.
- Long periods of rain can cause an increase of such fungal diseases as scab and leaf spots.

The more common diseases of landscape plants are described below.

Vascular Wilt

Vascular wilt fungi of shade trees are of two types:

- those that infect roots (Verticillium wilt), and
- those that infect stems (Dutch elm disease).

The organism that causes Verticillium wilt may be present in the soil. It spreads upward from the roots through sapwood and interferes with water movement and other plant functions. Dutch elm disease is transmitted by elm bark beetles.

In both diseases, leaf wilting, browning between veins, and leaf drop usually begin in one branch and progress through the tree. Dead and dying branches, sparseness of the crown, and reduced twig growth are common symptoms. Another is a discolored streaking in the wood of affected branches or in the main trunk.

Leaf Spots

Fungal leaf spots occur on many kinds of ornamental plants. They may begin as dark brown, pinhead-sized spots which sometimes have a yellow halo. Spots may enlarge to cover an entire leaf. As the spots become more abundant, leaves may yellow, die, and drop.

Leaf spots are more common in the early spring and fall. Wet conditions usually are necessary for infection. Healthy plants become infected when the fungus spores are:

- splashed onto them from infected leaves on the ground,
- blown to them by the wind, or
- carried to them on clothing and tools.

Scab

Apples, crab apples, pyracantha, and some other plants are susceptible to the scab fungi.

Symptoms include:

- spots on leaves and fruit, and
- premature defoliation

Scab first appears as olive-green spots on the new leaves. These spots become brown and velvety; then leaves turn yellow and drop prematurely. Fruit may become infected at any time with circular, olive-green spots that later become brown or black. The fungi overwinter in infected leaves and produce spores in the spring.

Powdery Mildew

Powdery mildew occurs on plants both in greenhouses and outdoors. Common hosts are rose,

zinnia, crab apple, euonymus, and crape myrtle. Powdery mildew may produce a white powdery coating on the leaves, buds, or stems of highly susceptible plants. The new growth is stunted and curled, and leaves may become dry and drop. The flower buds are often deformed and may fail to open properly.

Bacterial Fire Blight

Certain varieties of apple, flowering crab, pear, pyracantha, mountain ash, and quince are highly susceptible to fire blight. Hawthorn, rose, cotoneaster, spirea, and amelanchier are affected less seriously.

The signs of fire blight are:

- Blossoms and leaves suddenly wilt, turn dark brown, shrivel, and die, but usually remain attached.
- Secondary infections start in the small twigs, progress down the stem, and may involve whole branches.
- Blighted terminals may bend to look like a shepherd's crook.
- Dark streaking of the wood extends several inches beyond the diseased area.
- Cankers on limbs are shrunken, and are dark brown to purple. An orange gum or slime often oozes from them.

The bacteria overwinter in cankers on the plant. They are spread by:

- wind-blown rain,
- insects, and
- pruning tools.

Nematodes

Many nematodes live in the soil and feed on plant roots. Some kinds cause small knots on roots; others kill the tips of feeder roots.

The above-ground symptoms of nematode damage may include:

- yellowing of foliage,
- stunting, and
- a general decline of the plant.

It is difficult to distinguish between the symptoms of nematode damage and root rot infection. You may need to have soil and plant samples examined

in a laboratory to confirm a nematode infestation. Root-knot of boxwood is an example of a nematode disease of ornamentals.

Disease Control

Types of chemicals available for disease control include:

- Protective chemicals applied to foliage, flowers, and fruit. They are subject to weathering and must be reapplied regularly.
- Systemic chemicals. These can be applied less frequently.
- Soil fumigants. Use of these to control soil-borne fungi, bacteria, and nematodes is economically feasible in the production and establishment of high-value ornamentals.

There are no known chemicals for control of virus infections. With few exceptions, disease-controlling pesticides will not eradicate disease-producing agents after infection has occurred. Careful management, including pruning out of dead and dying plant parts and the removal of infected leaves, coupled with preventive use of the correct fungicides and bactericides, will prevent further spread. In some situations, routine preventive use of a pesticide is the only practical way to protect highly susceptible plants. Your local farm advisor can help you identify your pest problems and select the correct pesticide.

WEEDS

Many kinds of weeds are pests in landscape plantings.

Annual Weeds

Annual weeds are most troublesome in intensively cultivated ornamentals. Common annual weeds in ornamentals include:

- grasses (crabgrass, foxtailgrass, fall panicum, and barnyardgrass) which germinate during the spring and summer,
- annual bluegrass and annual brome grass, which germinate during the late summer or fall,
- annual broadleaf weeds (purslane, pigweed, and lambsquarters) which germinate during the warm season and are killed by hard frost, and
- those that survive freezing temperatures (horseweed, common chickweed, bittercress, and pepperweed).

Biennial and Perennial Weeds

Biennial and perennial weeds are most troublesome in uncultivated ornamentals. They have underground plant parts that survive from year to year. They are spread in several ways:

- Many spread easily when carried in soil, in root balls, and on cultivating equipment, as well as by seeds. These include Bermuda grass, Johnson grass, quackgrass, nutsedge, mugwort, and wild garlic.
- Seeds of perennial weeds such as dandelion and goldenrod are spread primarily by wind and water.
- Horsetail rush is spread by underground plant parts and by spores.

Weed Control

Consider both the weeds and the ornamental plants when choosing control methods. You can use cultural methods, mechanical methods, herbicides, or combinations of the three. Many weeds are resistant to some cultural or chemical controls. No herbicide is safe for all ornamental plants. Newly planted ornamentals usually are more easily injured by herbicides than established plantings. The label will tell you how to use a herbicide safely and effectively.

Herbicides kill weeds through the leaves or the roots or both. Selective herbicides kill some plants without killing others. Nonselective herbicides kill most plants in the area of application.

The main types of herbicides used in or around ornamentals are:

- preemergence herbicides,
- postemergence herbicides, and
- soil fumigants and sterilants.

Persistence varies with the herbicide and the dosage. Persistent herbicides may leave residues that may injure a sensitive crop planted later. Repeated applications of persistent herbicides also can injure ornamental plants under certain soil and climatic conditions. Granular formulations are an efficient way to apply preemergence herbicides. Postemergence herbicides usually are less persistent than preemergence herbicides. They usually must be applied as a directed spray.

Soil fumigants are nonselective and cannot be used in the root zones of desirable plants. Use fumi-

gants before planting. The label will specify waiting periods between treatment and planting.

Soil sterilants will control most weeds for long periods of time. In humid regions, however, no material is completely effective for more than one season. Soil sterilants are nonselective. They can damage nearby trees, shrubs, and turfgrass through root uptake or movement of the chemical by wind or water.

INSECTS AND MITES

Ornamental plants are damaged by many kinds of insects and mites. Some suck sap from plants, others chew on or tunnel in plant parts or cause damage in other ways.

Some plants are very susceptible to insects and mites and require intensive pest control. Other plants are rarely attacked by insects or mites. Pest insect infestations vary from year to year, and control is not always needed.

There are several categories of insects and mites that you should recognize. They can be grouped according to the part of the plant they feed on and the kind of injury they cause.

Insects and Mites that Damage Leaves, Buds, Fruits, and Flowers

CATERPILLARS are the larvae of butterflies or moths. Caterpillars chew plant parts and may completely defoliate a plant. Some form webs or tents on the branches. A few bore into the plant and feed inside. Some have more than one generation per year.

BEETLES are hardshelled insects. Many have spots, stripes, or other markings. Both adults and larvae may damage plant parts by boring into or chewing them. Some beetles are active only at night.

LEAFMINERS are the larvae of certain small flies, moths, or beetles. They feed inside the leaf. Damage appears as brown or discolored blotches or winding trails on the leaf. There may be more than one generation per year.

APHIDS are small, soft-bodied insects that suck sap through tiny needle-like mouthparts. There may be several generations in a single season. They may be green, red, or black. They feed on stems, terminals, or undersides of leaves.

Foliage often curls or is otherwise distorted. Some aphids transmit plant disease. Aphids produce honeydew, a sweet liquid which collects on the foliage. A black sooty mold may grow on the honeydew. Sooty mold is controlled by controlling the aphids.

MITES are closely related to insects. They are hard to see without magnification. Eggs, young, and adults all may be present on an infested plant at the same time. Some form webs on the lower leaf surface. Mites damage leaves by sucking sap. The foliage becomes stippled and may turn off-green, yellow, or orange. Mites may produce several generations in a single season.

THRIPS are tiny, sliver-like insects often less than $1/16$ inch in length. Most feed inside flowers, where flecking of blossoms results. Others live in the growing points of plants and cause distortion or bleaching of leaves.

Insect Pests of Trunks, Stems, or Branches

SCALE INSECTS AND MEALYBUGS may kill large branches or whole plants. Some attack leaves and buds. Both insects secrete a protective waxy substance which covers them.

Mealybugs move on the plants as both young and adults. Newly hatched scale insects (crawlers) move around on the plant. Mature scale insects, however, are securely fastened to the plant surface. They may be circular, oval, or pear-shaped. Large numbers may form crusts on the plant. They lay eggs underneath the protective covering.

BORERS are larvae of some moths and beetles. They do the most damage in the tissue just under the bark. Plants in poor health are more susceptible to attack by borers. One to several years may be required to complete a life cycle.

Insect Pests of Roots

GRUBS are the larvae of beetles or weevils. They usually are white with brown heads. Some have legs; others are legless. Grubs eat plant roots and may weaken or kill a plant.

ROOT BORERS are the larvae of moths or beetles. They are shaped like grubs or caterpillars. They are usually a whitish color. Root borers damage plants

by eating or hollowing out plant roots and crowns.

ROOT-FEEDING APHIDS weaken the root systems of plants. They damage roots by sucking sap, which may cause galls to form. Root aphids look like foliar-feeding aphids. Some are serious pests of foliage as well as roots.

GALLS are swellings of plant parts. Many kinds of insects and mites cause galls and live inside them. Galls are unsightly but usually are not harmful to the health of a plant.

Insect and Mite Control

To control insects and mites, direct the pesticide at the stage of the insect or mite that is causing the damage. In some situations, preventive use of insecticides may be necessary to protect plants from infestations. Your decision to use a preventive insecticide should be based on a previous history of infestation in your area. Your local extension agent can help to identify your pest problems and select the correct pesticide.

VERTEBRATE PESTS

Vertebrate animals may damage ornamentals in several ways. Some (such as mice and moles) feed on roots and crowns. Others (including mice, rabbits, deer, and woodpeckers) feed on stems, trunks, twigs, or foliage.

Barriers, trapping, repellents, and pesticides all help control vertebrate pests. Control of the insects on which vertebrate pests feed is essential.

TURFGRASS

INTRODUCTION

Pest control in turfgrass includes:

- good cultural practices, and, when necessary,
- chemical pest control.

Turfgrass problems often result from causes other than pests. These other causes include:

- improper watering,
- improper fertilization practices,
- injury from pesticides,
- accumulation of excessive thatch,
- improper selection of turfgrass species,
- improper mowing height,
- poor root systems,
- detrimental weather conditions.

- soil that is either too acid or too alkaline, or
- an accumulation of soluble salts in the soil.

Be sure to consider these factors when diagnosing and treating turfgrass problems.

DISEASE AGENTS

The major diseases of turfgrass are caused by:

- fungi, which can cause root rots and foliar diseases, and
- nematodes, which feed on the roots.

Stands of diseased grass may look thin and unthrifty or contain streaks or circular patches of dead grass. Some of the more common diseases of turfgrasses are:

Helminthosporium

Helminthosporium causes leaf spots and crown rots. Spots on the leaves usually begin as small purplish, reddish-brown areas about the size of a pinhead. These enlarge to form tan to light-brown spots with reddish-brown margins. When the disease is severe, the spots girdle the leaves at the base and cause them to yellow. A severe infection may cause a general fading out of turfgrass. Helminthosporium diseases are more severe during long periods of wet weather.

Rusts

Rusts form elongated, reddish pustules containing spores which appear on stems, leaves and leaf sheaths of grasses. The reddish spores adhere to the fingers when the pustules are rubbed. Rusts develop rapidly during moderately warm, moist weather. Moisture in the form of dew for 10 to 12 hours is sufficient for the spores to infect plants.

Pythium Blights

Pythium blights are among the most destructive turfgrass diseases. Grasses most commonly affected are bentgrasses, Bermuda grasses, fescues, and rye grasses. The disease is first seen as small, irregularly-shaped, watersoaked, greasy patches ½ to 4 inches in diameter. A cottony growth may be present early in the morning. Diseased areas may eventually range from 1 to 10 feet in diameter. Pythium blights develop best in warm, humid weather. They

are most severe on grass grown under high nitrogen fertilization.

Rhizoctonia Brown Patch

Under conditions of close mowing, Rhizoctonia brown patch appears as irregularly shaped patches of blighted turfgrass that range in size from a few inches to 2 feet or larger. At first, the patches are purple-green in color. They then fade to a light brown. When the grass is wet, the diseased patches frequently have dark, purplish margins (smoke rings).

When high mowing is practiced, the leaves wither and rapidly fade to a light brown. The patches may be irregular and range up to 50 feet in diameter. Rhizoctonia brown patch develops best during long periods of humid weather. The disease usually occurs during hot weather (80–90 degrees F). Grass grown under high nitrogen fertilization is more susceptible to the disease.

Fusarium Patch

Fusarium fungus causes roughly circular patches as large as 12 inches in diameter in grasses. The grass leaves first become water-soaked. Later they turn reddish-brown, then become bleached. White or pinkish fungus threads may be seen in early morning. The disease is most prevalent during cool, moist periods such as the prolonged winter rainy season.

Slime Molds

Slime molds appear as dull-gray to light-blue masses of powdery growth on the surfaces of the leaves. They are most common during long periods of light rainfall. Although they are unsightly, they do not damage the grass. Controls are not necessary.

Fusarium Blight

Fusarium blight causes brown patches 1 to 3 feet in diameter. The patches are similar to those caused by other turfgrass diseases, but they have green tufts (frog eyes) in the center. Fusarium blight is most severe during periods of high day and night temperatures. Lush grass with an accumulation of 1 inch or more of thatch is highly susceptible to severe outbreaks.

Fairy Rings

Fairy rings are seen as circles of darker green, faster-growing turfgrass ranging from 2 to several hundred feet in diameter. They are often surrounded by mushrooms, toadstools, or puffballs. These fungi may prevent water from penetrating the soil.

Nematodes

Many kinds of nematodes feed on the roots of turfgrasses and reduce their vigor. Nematode injury may be confused with nutritional problems, insufficient water, compact soil, or any other factor which restricts root development.

Symptoms of nematode injury include:

- thinning or completely killed areas,
- pale green to yellow color,
- excessive wilting, and
- poor response to fertilization.

The best way to identify nematode problems is with a laboratory examination of soil or plants.

Disease Controls

All turfgrass diseases are easier to prevent than to cure. To minimize the possibility of disease, plant the right grass variety for a particular climate at the recommended rate and time of year. Weakened, nonadapted grasses are more susceptible than normal to certain turf-attacking fungi, and to such stresses as drought and hot, dry winds.

Recommended cultural practices, such as proper mowing, adequate fertilization, judicious and timely irrigation, and frequent aerification, help prevent diseases by maintaining a vigorously growing turf. In general, a properly maintained turfgrass is less severely damaged by diseases and is able to recover more quickly than a poorly maintained one, because its ability to form new shoots and roots to replace those damaged or killed by diseases is not impaired.

One cultural practice important to disease prevention is fertilization, specifically the amount of nitrogen applied. Too much nitrogen can result in a soft, lush growth of the grass — a condition favorable to some diseases. On the other hand, turf deficient in nitrogen is susceptible to certain other diseases. If cultural

methods do not control diseases, using certain effective fungicides is recommended. Fungicides are usually most effective if applied before a disease becomes severe. However, if the disease does become severe, it is usually necessary to apply the fungicide more frequently and at a higher rate.

WEEDS

Any plant can be considered a weed if it is growing where it is not wanted. Bentgrass, for example, would be a weed in a bluegrass lawn. To plan a good weed control program, you must:

- identify the desirable turfgrass,
- identify the existing weeds, and
- know what other weeds are likely to become a problem.

Annual Weeds

Annual weeds complete their life cycle in less than one year. Because climatic conditions influence the timing of the life cycle, the correct time for control varies from place to place, year to year, and from one species to another. It is often desirable to establish turfgrass in the fall so the freezing weather will control summer annual weeds. In established turfgrass, the chemical control of summer annual weeds after midsummer may not be necessary or desirable.

SUMMER ANNUAL WEEDS common to turfgrass are:

Broadleaf Weeds

henbit
knotweed
spurge

Grass Weeds

crabgrass
goosegrass
barnyardgrass
foxtailgrass
stinkgrass

WINTER ANNUALS are common in new turfgrass. After the first year, good management and dense turfgrass usually provide satisfactory control. Examples are:

Broadleaf

common chickweed
shepherdspurse

Grass Weeds

cheat

Biennial Weeds

Biennial weeds normally occur at the same time

as perennial broadleaf weeds. Controls are similar. Examples are: roundleaf mallow and wild carrot.

Perennial Weeds

Perennials, both broadleaf and grasses, occur widely as turfgrass weeds. Examples are:

Broadleaf Weeds

dandelion
wild garlic
dichondra
plantain
mouse-ear chickweed
red sorrel

Grass Weeds

Bermuda grass
bentgrass
tall fescue
quackgrass
nimblewill
torpedograss
nutsedge

Weed Control

The presence of weeds in turfgrass does not always require the use of herbicides. In areas that contain sensitive plants, it may be better to avoid the use of herbicides than to risk injury. In some locations, any kind of plant cover may be better than dead plants or bare ground.

Granular formulations are effective for pre-emergence herbicides. Sprays are better for post-emergence control where foliar coverage is needed.

BROADLEAF WEEDS—Several postemergence herbicides are used to selectively control annual, biennial, and perennial broadleaf weeds in turfgrasses. They can be used alone or as combinations of more than one active ingredient. Spring and fall applications of postemergence herbicides normally give satisfactory control and reduce the possibility of damage to nontarget plants. Young weeds are usually more susceptible to herbicides. Spot treatments are best for scattered weed populations. Weather conditions affect control results.

GRASS WEEDS—Control of annual grasses is best achieved with:

- preemergence herbicides for general infestations,
- spot treatment with postemergence herbicides for localized infestations.

Few herbicides are safe for use on newly seeded turfgrass. Some preemergence herbicides applied in the spring adversely affect germination of turfgrasses seeded in the fall. Certain varieties of turfgrasses are more prone to injury by some herbicides. Check labels for precautions.

Perennial grass weeds are the most difficult to control. No herbicides are available which will control these weeds without damaging cool season turfgrass. Some will selectively control them in warm season turfgrass. Soil fumigants and nonselective herbicides are sometimes used.

INSECTS

When examining turfgrass for insects, look for:

- thinned grass stands,
- dying or dead patches,
- discolored or withered blades,
- chewed or frayed blades,
- frass or webbing,
- small holes, mounds, or burrows, or
- presence of large numbers of bird and animal droppings.

Some of the more troublesome turfgrass insect pests are:

Sod Webworms

Sod webworm caterpillars are 1 inch or less in length. They are off-white with parallel rows of small dark spots. The adults are cigar-shaped, buff-colored moths. The caterpillars chew off grass stems and leaves above the soil line. Damage shows up as small dead spots. When many sod webworms are present, the spots join to form large, irregularly shaped brown patches. Adult sod webworms do not damage turfgrasses.

Billbugs

Billbugs are small, dark-colored beetles with snouts. Adults lay eggs in turfgrass stems in late spring. The eggs hatch into legless larvae. The larvae eat their way down the stems and into the crowns. Adults feed on leaves and stems, but cause less damage than the larvae. Damage shows up in late summer as small dead patches of turfgrass. Damaged plants break off at the crown if pulled on.

Insect Control

Insects that attack turfgrass at or below the soil surface can be controlled only by directing the pesticide at the soil surface and watering it in to contact the pests. Foliage-feeding insects can be controlled by directing the insecticide at the turfgrass foliage.

Watering in an application directed at foliage

feeders will move the insecticide below the area where the insect pest is feeding and the desired control will be lost. In some areas, preventive applications of insecticides will minimize damage from soil insect pests. More than one pest may be causing damage at the same time. Each may require different timing and placement of insecticide for control. Be sure to consider this when you develop a treatment schedule. Your local extension agent can help you identify pest problems and select the correct pesticide.

VERTEBRATE PESTS

Vertebrate animals may damage large areas of turfgrass while they are searching for grubs or other soil-infesting insects. They include:

- mice,
- voles,
- skunks,
- moles,
- raccoons,
- foxes,
- squirrels, and
- birds.

Control of turfgrass-damaging insects also helps control damage by vertebrate animals, because it reduces their food supply.

For additional information on vertebrates see applicator training guide: "Vertebrate Pest Control".

PHYTOTOXICITY

Phytotoxicity is undesirable injury to plants. Symptoms of phytotoxicity include:

- leaf drop,
- stunting,
- overgrowth,
- discolored foliage,
- leaf curl, and
- stem distortion.

The cause of phytotoxicity may be easy to determine or it may be subtle and hidden. Pesticides can cause phytotoxicity. Other causes that create similar symptoms are:

- insects and disease agents,
- insufficient moisture,
- improper fertilization, and

- other adverse growing conditions.

Factors that may contribute to pesticide phytotoxicity include:

- high air temperature during and immediately after pesticide application,
- excessive rates of pesticide application,
- too little water,
- uneven distribution of pesticide,
- mixing liquids or emulsifiable concentrates with wettable powders,
- mixing fertilizers with pesticides,
- variety and species differences.

Take special care to avoid injury to landscape plants and turfgrass when using herbicides. Some herbicides leave residues in spray tanks that will injure desirable plants. Use separate sprayers for herbicides.

ENVIRONMENTAL CONCERNS

To control drift and vaporization:

- Apply pesticides when wind speeds are low.
- Use lowest practical operating pressure and largest practical nozzle opening.
- Keep nozzle as close to target as possible.
- Avoid using airblast sprayers and dusters when working near sensitive plants and areas inhabited by animals.
- Do not apply herbicides with airblast sprayers.
- When possible, select products with low volatility.

To control the adverse effects of pesticide movement:

- Use special precautions when using pesticides on slopes.
- Select the least hazardous pesticide that will do the job.
- Use the lowest effective rate of application.
- If possible, maintain a buffer zone between the area to be treated and sensitive areas.
- Use mulches.
- Consider the chances of heavy rainfall.
- Regulate the amount and duration of irrigation.
- Be aware of the potential for ground water contamination.
- Avoid carrying treated material or the pesticide residue from the target area to other areas.

You must know the persistence of pesticides you apply to ornamentals and turfgrass, especially where:

- adjacent areas may be affected,
- treated soil is used to grow other plants, or
- humans, pets, or other animals are present.

Repeated applications of some pesticides to the same area may cause harmful residues

PROTECTING ANIMALS AND PEOPLE

Keep animals and people away during application and until spray has dried or dust has settled. Keep them away from areas of potential drift and runoff. Remove toys, pet food dishes, birdfeeders, and other articles from the site before applying a pesticide. Do not use pesticides when people or pets cannot be excluded during the reentry period specified on the label.

APPLICATION

Methods of application vary with:

- the kind of pesticide,
- the host, and
- the target pest.

Application equipment must be able to deliver a thorough coverage of the correct amount of pesticide to the plant parts which need protection.

Low-pressure, low-volume sprayers or granular applicators can be used for control of:

- soil or foliage pests of ornamentals,
- diseases or insects on turfgrass, or
- weeds.


High-pressure hydraulic or airblast sprayers are not often used on ornamentals or turfgrass. You can use them for spraying large trees.

AREA MEASUREMENTS

To determine how much pesticide you will need to do a job, you must measure the area to be treated. If the area is a rectangle, circle, or triangle, simple formulas may be used.


Rectangles: The area of a rectangle is found by multiplying the length by the width.

Area = Length $\times$ Width.




Circles: The area of a circle is the radius (one-half the diameter) squared and then multiplied by 3.14.

Area = 3.14 $\times$ the radius squared.




Triangles: The area of a triangle is one-half the base multiplied by the height.

$$\text{Area} = \frac{b \times h}{2}$$



Irregularly shaped turfgrass areas often can be reduced to one or more of these common shapes. Calculate the area of each and add them together to obtain the total area.

Example:



Area A + B + C = Total Area

Another way is to establish a line down the middle of the property for the length, and then measure from side to side at several points along this line. Areas with very irregular shape require more side to side measurements. The average of the side measurements can be used as the width. The area is then calculated as a rectangle.

Area = Length $\times$ Width. Example:

WEIGHTS AND MEASURES

Weights

1 ounce	= 28.35 grams
16 ounces	= 1 pound = 453.59 grams
1 gallon water	= 8.34 pounds = 3.785 liters = 3.78 kilograms

Liquid Measures

1 fluid ounce	= 2 tablespoons = 29.573 milliliters
16 fluid ounces	= 1 pint = 0.473 liter
2 pints	= 1 quart = 0.946 liter
8 pints = 4 quarts	= 1 gallon = 3.785 liters

Length

1 foot	= 30.48 centimeters
3 feet	= 1 yard = 0.9144 meter
16½ feet	= 1 rod = 5.029 meters
5,280 feet	= 320 rods = 1 mile = 1.6 kilometers

Area


1 square foot	= 929.03 cm ²
9 square feet	= 1 square yard = 0.836 square meter
43,560 square feet	= 160 square rods = 1 acre = 0.405 hectare

Speed

1.466 feet per second	= 88 feet per minute = 1 mph = 1.6 kilometers per hour (kph)
-----------------------	--

Volume

27 cubic feet	= 1 cubic yard = 0.765 cubic meter
1 cubic foot	= 7.5 gallons = 28.317 cubic decimeters




Length = line AB

$$\text{Width} = \frac{\text{line C} + \text{D} + \text{E} + \text{F} + \text{G}}{5}$$

A third method is to convert the area into a circle. From a center point measure distance to the edge of the area in 10 to 20 increments. Average these measurements to find the average radius. Then calculate the area, using the formula for a circle. Area = $3.14 \times \text{the radius squared}$.

Example:



$$\text{Area} = (3.14) \times \left(\frac{\text{line A} + \text{B} + \text{C} + \text{D} + \text{E} + \text{F} + \text{G} + \text{H} + \text{I} + \text{J} + \text{K} + \text{L}}{12} \right)^2$$